



Methanol masers and millimetre lines:  
a common origin in protostellar envelopes

Cover: Photograph of a shot glass filled with Rhodamine 6G laser dye dissolved in methanol. The dye is illuminated by the 355 nm output of a pulsed Nd:YAG laser. The exposure time of 25 s corresponds to ~250 pulses, each with a duration of 4 ns. Produced with a lot of help from Dr. Guss.

# Methanol masers and millimetre lines: a common origin in protostellar envelopes

PROEFSCHRIFT

ter verkrijging van  
de graad van Doctor aan de Universiteit Leiden,  
op gezag van de Rector Magnificus prof.mr. P.F. van der Heijden,  
volgens besluit van het College voor Promoties  
te verdedigen op dinsdag 6 december 2011  
klokke 11.15 uur

door

Karl Johan Erik Torstensson  
geboren te Högsbo, Sweden  
in 1977

## Promotiecommissie

Promotores: Prof. dr. E. F. van Dishoeck  
Co-promotor: Dr. H. J. van Langevelde

Overige Leden: Prof. dr. K. Kuijken  
Prof. dr. A. G. G. M. Tielens  
Prof. dr. J. E. Conway  
Dr. F. F. S. van der Tak  
Dr. W. H. T. Vlemmings

Onsala Space Observatory  
SRON Netherlands Institute for Space Research  
Chalmers University of Technology

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Star formation . . . . .	1
1.2	Molecular astrophysics . . . . .	3
1.3	Masers . . . . .	6
1.4	Radio interferometry . . . . .	8
1.5	This thesis . . . . .	8
1.6	Main conclusions . . . . .	10
1.7	Future prospects and outlook . . . . .	11
<b>2</b>	<b>Dynamics of the methanol masers around Cepheus A HW2</b>	<b>13</b>
2.1	Introduction . . . . .	14
2.2	Observations and data reduction . . . . .	15
2.2.1	6.7 GHz data . . . . .	15
2.2.2	12.2 GHz data . . . . .	17
2.3	Results . . . . .	17
2.3.1	6.7 GHz results . . . . .	17
2.3.2	12.2 GHz results . . . . .	22
2.4	Analysis . . . . .	24
2.4.1	Ring model . . . . .	24
2.4.2	Parallax - distance . . . . .	26
2.5	Discussion . . . . .	27
2.6	Conclusions . . . . .	28
<b>3</b>	<b>Distribution and excitation of thermal methanol in Cepheus A</b>	<b>31</b>
3.1	Introduction . . . . .	33
3.2	Observations and data reduction . . . . .	35
3.3	Results . . . . .	37
3.3.1	Methanol lines . . . . .	37
3.3.2	Spatial distribution of methanol . . . . .	39
3.4	Analysis . . . . .	41

## Contents

3.4.1	Rotation diagrams . . . . .	41
3.4.2	Population diagram modelling . . . . .	43
3.4.3	Spatial distribution of the excitation . . . . .	43
3.4.4	Non-LTE analysis . . . . .	45
3.4.5	H <sub>2</sub> column density . . . . .	48
3.4.6	Methanol abundance . . . . .	48
3.5	Discussion . . . . .	50
3.5.1	Outflow morphology . . . . .	50
3.5.2	Methanol distribution . . . . .	50
3.5.3	Physical conditions . . . . .	51
3.5.4	Origin of maser emission . . . . .	52
<b>4</b>	<b>Thermal methanol toward 6.7 GHz methanol maser sources</b>	<b>55</b>
4.1	Introduction . . . . .	57
4.2	Observations and data reduction . . . . .	58
4.3	Results . . . . .	61
4.3.1	AFGL 5142 . . . . .	61
4.3.2	DR21 (FIR1 & FIR2) . . . . .	64
4.3.3	G23.207−00.377 . . . . .	67
4.3.4	G23.389+00.185 . . . . .	70
4.3.5	G23.657−00.127 . . . . .	72
4.3.6	G24.541+00.312 . . . . .	73
4.3.7	G40.62−0.14 . . . . .	74
4.3.8	G73.06+1.80 . . . . .	76
4.3.9	G78.12+3.63 . . . . .	78
4.3.10	L1206 . . . . .	80
4.3.11	S255 . . . . .	82
4.3.12	W3(OH) . . . . .	85
4.4	Analysis . . . . .	88
4.4.1	Rotation diagram analysis . . . . .	88
4.4.2	CH <sub>3</sub> OH gas distribution . . . . .	89
4.4.3	Population diagram analysis . . . . .	91
4.5	Discussion . . . . .	92
4.5.1	Excitation of CH <sub>3</sub> OH gas . . . . .	92
4.5.2	Morphology . . . . .	94
4.6	Conclusions . . . . .	94
<b>5</b>	<b>Dynamics of 6.7 GHz methanol masers in high-mass star-forming regions</b>	<b>97</b>
5.1	Introduction . . . . .	99
5.2	Observations and data reduction . . . . .	100
5.3	Results . . . . .	101
5.4	Discussion . . . . .	104
5.5	Summary . . . . .	108

## Contents

<b>Bibliography</b>	<b>113</b>
<b>Nederlandse Samenvatting</b>	<b>119</b>
<b>Publications</b>	<b>125</b>
<b>Curriculum Vitae</b>	<b>127</b>
<b>Acknowledgements</b>	<b>129</b>



